

Just Accepted by Nanotoxicology

Experimental evidence of false positive Comet test results due to TiO₂ particle - assay interactions

Rajapakse Katarina*, Drobne Damjana †, ‡, §, Kastelec Damijana¶, Marinsek-Logar Romana*

doi:10.3109/17435390.2012.696735

Abstract

We have studied the genotoxicity of TiO₂ particles with a Comet assay on a unicellular organism, *Tetrahymena thermophila*. Exposure to bulk or nano-TiO₂ of free cells, cells embedded in gel or nuclei embedded in gel, all resulted in a positive Comet assay result but this outcome could not be confirmed by cytotoxicity measures such as lipid peroxidation, elevated reactive oxygen species or cell membrane composition. Published reports state that in the absence of cytotoxicity, nano- and bulk TiO₂ genotoxicity do not occur directly and a possible explanation of our Comet assay results is that they are false positives resulting from *post-festum* exposure interactions between particles and DNA. We suggest that before Comet assay is used for nanoparticle genotoxicity testing, evidence for the possibility of *post-festum* exposure interactions should be considered. The *acellular* Comet test described in this report can be used for this purpose

© Informa UK, Ltd. This provisional PDF corresponds to the article as it appeared upon acceptance. Fully formatted PDF and full text (HTML) versions will be made available soon.

DISCLAIMER: The ideas and opinions expressed in the journal's *Just Accepted* articles do not necessarily reflect those of Informa Healthcare (the Publisher), the Editors or the journal. The Publisher does not assume any responsibility for any injury and/or damage to persons or property arising from or related to any use of the material contained in these articles. The reader is advised to check the appropriate medical literature and the product information currently provided by the manufacturer of each drug to be administered to verify the dosages, the method and duration of administration, and contraindications. It is the responsibility of the treating physician or other health care professional, relying on his or her independent experience and knowledge of the patient, to determine drug dosages and the best treatment for the patient. *Just Accepted* articles have undergone full scientific review but none of the additional editorial preparation, such as copyediting, typesetting, and proofreading, as have articles published in the traditional manner. There may, therefore, be errors in *Just Accepted* articles that will be corrected in the final print and final online version of the article. Any use of the *Just Accepted* articles is subject to the express understanding that the papers have not yet gone through the full quality control process prior to publication.

Experimental evidence of false positive Comet test results due to TiO₂ particle - assay interactions

Rajapakse Katarina^{*}, Drobne Damjana^{†, □, §}, Kastelec Damijana[¤], Marinsek-Logar Romana^{*}

^{*} Department of Animal Sciences, Biotechnical faculty, University of Ljubljana, Groblje 3, SI-1230 Domzale, Slovenia Fax: +38617241005; E-mail: katarina.ales@bf.uni-lj.si, romana.marinsek@bf.uni-lj.si

[†] Department of Biology, Biotechnical faculty, University of Ljubljana, Vecna pot 111, SI-1000 Ljubljana, Slovenia; E-mail: damjana.drobne@bf.uni-lj.si

[□] Centre of Excellence in Advanced Materials and Technologies for the Future (CO NAMASTE), Jamova 39, SI-1000 Ljubljana, Slovenia;

[§] Centre of Excellence in Nanoscience and Nanotechnology (CO Nanocenter), Jamova 39, SI-1000 Ljubljana, Slovenia; E-mail:

[¤] Department of Agriculture, Biotechnical faculty, University of Ljubljana, Vecna pot 101, SI-1000 Ljubljana, Slovenia E-mail: damijana.kastelec@bf.uni-lj.si

Address correspondence to:

Damjana Drobne: Department of Biology, Biotechnical faculty, University of Ljubljana, Vecna pot 111, SI-1000 Ljubljana, Slovenia; Phone number: + 386(1) 320 33 75, E-mail: damjana.drobne@bf.uni-lj.si

Key words: *Tetrahymena thermophila*, DNA damage, nanoparticles, nanotoxicity

ABSTRACT

We have studied the genotoxicity of TiO₂ particles with a Comet assay on a unicellular organism, *Tetrahymena thermophila*. Exposure to bulk or nano-TiO₂ of free cells, cells embedded in gel or nuclei embedded in gel, all resulted in a positive Comet assay result but this outcome could not be confirmed by

cytotoxicity measures such as lipid peroxidation, elevated reactive oxygen species or cell membrane composition. Published reports state that in the absence of cytotoxicity, nano- and bulk TiO₂ genotoxicity do not occur directly and a possible explanation of our Comet assay results is that they are false positives resulting from *post-festum* exposure interactions between particles and DNA. We suggest that before Comet assay is used for nanoparticle genotoxicity testing, evidence for the possibility of *post-festum* exposure interactions should be considered. The *acellular* Comet test described in this report can be used for this purpose.

Running head: False positive genotoxicity of NPs due to *post-festum* exposure interactions

JUST ACCEPTED

INTRODUCTION

Genotoxicity has been defined by the International Conference of Harmonization in an ICH-Guideline as deleterious change in the genetic material induced by any mechanism. Damage to DNA results in cellular dysfunction and may therefore initiate and promote mutagenesis and carcinogenesis, or impact fertility (Sathya et al., 2010). Because of this, data on genotoxicity are of great importance in regulatory health risk assessment.

Genotoxicity of nanoparticles (NP) has frequently been documented (Sathya et al., 2010) (Landsiedel et al., 2010, Karlsson, 2010) and the mechanisms of this genotoxicity include direct primary genotoxicity driven by direct interaction of NPs with DNA (Donaldson et al., 2010) and indirect primary genotoxicity resulting from oxidative stress (Nel et al., 2006). The latter Oxidative stress occurs when NPs are transported into the nucleus (Chen and von Mikecz, 2005) (AshaRani et al., 2009) or when the nuclear membrane breaks down during mitosis (Karlsson, 2010). An example of an indirect mechanism is enhancement of the permeability of the lysosomal membrane, leading to release of DNase which, transported to the nucleus, can degrade DNA (Banasik et al., 2005). Secondary indirect mechanisms of nanoparticle genotoxicity are associated with inflammation (Trouiller et al., 2009).

With the advent of nanotechnology, it is essential to define a reliable test system with which to assess the genotoxic potential of engineered NPs (Warheit and Donner, 2010, Gonzalez et al., 2011). Guidelines provided by the Organisation for Economic Cooperation and Development (OECD) include *in vitro* genotoxicity testing, but these tests are designed basically for water-soluble chemicals and so may not be suited to testing of the genotoxicity of NPs. Nanoparticles interfere with test media, modifying the biological potential of the NPs and they may also interact with the test system, affecting the test results (Sathya et al., 2010); (Greim and Norppa, 2010). In an attempt to clarify this issue, the OECD has established projects designed to evaluate the relevance and reproducibility of genotoxicity assays (see (Warheit and Donner, 2010). Stone et al. (2009) have shown the importance when assessing direct primary genotoxicity of accurate distinction of artefacts and the possible interaction of test components with

nanoparticles remaining in the test system after exposure. Residual NPs may come into contact with nuclear DNA during tests affecting the test and this may also happen when NPs present inside cells in the cellular lysosomes or food vacuoles are released during the tests.

In recent research of genotoxicity and nanoparticles, the Comet assay has been one of the most frequently used tests (Landsiedel et al., 2009) (Karlsson, 2010). Recently, Karlsson (2010) reviewed 46 papers dealing with the genotoxicity of NPs by the Comet assay, and concluded that majority of the NPs tested caused DNA strand breaks. However, the possibility of interaction of NPs with the chemicals used in the assay was cited and the use of additional methods, distinct from the Comet assay was suggested for the measurement of DNA damage. Further mutagenicity studies have also been recommended. Landsiedel et al. (2009) suggested use of a battery of standardized genotoxicity tests covering a wide variety of potential mechanisms and suggested that at least two genotoxicity tests should always be implemented.

At present, there are four techniques in common use for *in vitro* testing the genotoxicity of nanoparticles. These are the Ames test, the Chromosomal Aberration Test, the Comet assay and the Micronucleus test. Of these, the Comet assay is the most popular because: (1) it is sensitive and capable of detecting low levels of DNA damage; (2) it requires only small numbers of cells per sample; (3) it is relatively inexpensive; and (4) it requires relatively small amounts of test substance (Tice et al., 2000). Among the limitations and disadvantages of the Comet assay are its failure to detect: (1) aneugenic effects; (2) epigenetic mechanisms of DNA damage (Dhawan et al., 2009); and (3) fixed mutations (Stone et al., 2009). There are also some serious obstacles to use of the Comet assay for NP genotoxicity studies. Karlsson (2010), for example has shown the presence of nanoparticles (nano-TiO₂ and nano-CuO), in heads of the comets in the gels, while intracellular localization of particles investigated by TEM did not reveal particles in cell nuclei. The possibility of post-exposure particle DNA interactions was also discussed by Lin et al. (Lin et al., 2009) who, studying the genotoxicity of Ge nanoparticles by the Comet assay. They noted a statistically higher level of DNA damage in exposed cells when compared with control cells and speculated that, since

nanoparticles of Ge readily adhere to cell surfaces, nanoparticles in or attached to the cells caused the damage during the assay process.

The features that motivated the selection of *T. thermophila* as a model organism for this study are: (1) It is a one-cell eukaryotic organism. Thus, the data obtained by Comet assay correspond to the impact of TiO₂ on whole organism DNA, and in summary the effects are measured on the genome of the entire cell population. (2) Its short generation time and its axenical culture are especially advantageous for studying genotoxicity. (3) As protists have highly developed systems for internalisation of nanoscale (100 nm or less) and microscale (100 – 100,000 nm) particles (Frankel, 2000) they are very good model organisms for nanotoxicology (Holbrook et al., 2008) (Kahru et al., 2008). (4) It has been used in toxicology for decades as a useful model organism for cellular and molecular biologists as well as for environmental research (Sauvant et al., 1999, Gutiérrez et al., 2003).

The aim of the present study was to provide experimental evidence on the possibility that NPs interact with the DNA *post festum*, during a Comet assay. We used an unicellular model organism *T. thermophila* to assess genotoxicity by a Comet assay and cytotoxicity by conventional markers. In our study, three exposure scenarios were used: (a) *in vivo* exposure; *T. thermophila* was incubated in a suspension of particles – both nanoparticles and bulk TiO₂; (b) *in vitro* exposure; *T. thermophila* was embedded in gels which were incubated in a suspension of particles; (c) *acellular* exposure; only nuclei were embedded in gels and the gels with embedded nuclei were incubated in a suspension of particles. We chose to examine nano-TiO₂ particles for which a substantial amount of genotoxicity data already exists. (Trouiller et al., 2009) suggested that DNA damage results not from direct primary effects of nano-TiO₂ but rather from ROS generation, and is therefore a primary indirect effect. Very same was confirmed also by Petkovič et al. (2011 a, b) (Petkovic et al., 2011a) (Petkovic et al., 2011b). Consequently, we hypothesize that *in vitro* exposure (cells embedded in gel) and *acellular* exposure of only nuclei to nano-TiO₂ would fail to produce a positive result in a Comet assay since ROS generation, a primary indirect effect, would be absent. If *in vitro* and *acellular* exposure were to lead to a positive Comet assay, this would suggest that particles could

damage DNA during the tests, producing the positive Comet test result. In such cases, the use of Comet assay would have to be critically reconsidered.

MATERIALS AND METHODS

Chemicals.

Unless otherwise specified, reagents were purchased from Sigma Aldrich Co (St. Louis, MO, USA), Merck (Darmstadt, Germany) or Biolife (Milan, Italy). TiO₂ nanoparticles with 99.7% purity were supplied in the form of a powder.

***T. thermophila* growth conditions.**

Axenic cultures of *T. thermophila* from the Protoxkit FTM (MicroBioTests Inc.) were grown for 24 h in the dark at 25 °C in a semidefined-proteose-peptone based “rich” medium (RM) (Schultz, 1997). The cell density obtained after incubation in these culture conditions was approximately 10⁵ cells/ml.

Exposure conditions.

The cells were harvested by 3 min centrifugation at 60 rcf. Cells were washed and resuspended in a “poor” medium (PM), which consisted of the semidefined-proteose-peptone based medium used by Schultz (Schultz, 1997), but lacking yeast extract and bacteriological peptone. The pH of the medium was adjusted to 7.4 and temperature was maintained at 25 °C for the entire experiment. All experiments were performed in 100 ml batch cultures that were maintained in Erlenmeyer flasks and aerated by shaking at 90 rpm in an incubator in the dark.

After 1 h in the PM, cells were exposed to bulk or nano-TiO₂. The final concentration of particles in the medium, either bulk or nano-, was 0.1 and 100 µg/ml. Following the addition of TiO₂, *T. thermophila* cultures were incubated at 25 °C for 4 h. For each concentration of bulk or nano-TiO₂, three independent

assays were carried out. A supplementary set of three replicates without TiO₂, was set up as a control for each assay. After 4 h treatment with TiO₂ bulk or nanoparticles, 15 ml of cell suspension was harvested for the purpose of cellular fatty acid composition analysis by gas chromatography.

Bulk and nano-TiO₂ tested suspension.

Aqueous dispersions of nanoparticles were put on carbon-coated grids, dried at room temperature, examined with a 200-keV field emission transmission-electron microscope (Philips CM 100; Koninklijke Philips Electronics, Eindhoven, The Netherlands), and analyzed by transmission-electron diffraction to identify the TiO₂ crystal phase.

Bulk TiO₂ and 15 nm TiO₂ nanoparticles were dispersed in PM before treating the cell cultures. Bath sonication for 30 min was used to disperse particle agglomerates in stock solutions.

The suspensions of nanoparticles (1000 µg/ml) were inspected by dynamic light scattering (DLS) using a 3D DLS-SLS (dynamic light scattering - static light scattering spectrometer: LS Instruments, Fribourg, Switzerland). This allows the assessment of hydrodynamic radii of particles in extremely turbid suspensions by a so-called 3D cross-correlation technique that eliminates multiple scattering of light. As the light source a HeNe laser operating at a wavelength of 632.8 nm was used and scattering was measured at an angle of 90°.

Zeta potentials of TiO₂ nanoparticle suspensions (1000 µg/ml) were measured with ZetaPals, (Brookhaven Instrument Corporation) in the PM medium, and were used to assess the exposure to living cells.

Assessment of cellular fatty acid composition by gas chromatography.

T. thermophila cells were harvested by centrifugation at 60 rcf for 10 min of 15 ml culture samples. The pellets were resuspended in sterile double distilled water (1 ml) then frozen at -20 °C and lyophilized. Lipids were transesterified using a HCl/MeOH procedure (Dionisi et al., 1999). Dried samples were pulverized and transferred to screw cap test tubes. First, the sample was mixed with hexane (0.5 ml). Then

1.5M HCl in MeOH (1 ml) and pure MeOH (1 ml) were added and the test tubes were filled with N₂ and incubated at 80 °C for 10 min. The reaction was stopped by cooling the tubes in ice. Following addition of double distilled water (2 ml), each reaction mixture was vigorously mixed for 1 min and centrifuged for 30s at 670 rcf. The organic phase was transferred to a vial under N₂ and the samples were stored at -20 °C prior to analysis.

Fatty acid methyl esters were separated by capillary gas chromatography using Omegawax TM 320 (30m x 0.32 mm ID x 0.25 mm) capillary column with polyethylene glycol as the stationary phase. The gas chromatography system used was an Agilent 6890 series GC equipped with Agilent 7683 Automatic Liquid Sampler, 7683 Injector and FID detector and helium as the carrier gas with a flow rate of 2.0 ml/min, split ratio 10:1. The initial temperature for analysis was 185 °C and the final temperature was 215 °C. The injected volume was 2 µl and the run time was 54 min. Fatty acid methyl esters were identified from their retention times and results were calculated using response factors derived from chromatographic standards of known composition (Nu Chek Prep, GLC-85, Nu-Chek Prep Inc., Elysian, MN, USA).

Results were analyzed using ChemStation Plus® software. Membrane fatty acids which were less than 0.5% of total fatty acids were designated as trace fatty acids and were not considered further. Statistical analysis of the compositional data was used to evaluate differences in average fatty acid composition between different treatments (size and concentration of particles). Multivariate analysis of variance on isometric log-ratio transformations of the composition data was carried out.

Assessment of the extent of lipid peroxidation by quantitation of malondialdehyde.

Lipid peroxidation was tracked by the formation of malondialdehyde (MDA), a lipid peroxidation by-product that reacts with thiobarbituric acid (Ortega-Villasante et al., 2005). An aliquot of the culture (15 ml) was harvested by centrifugation at 6700 rcf for 10 min. Cells were homogenised by sonication for 3 min in an ice-cold water bath. To measure total protein concentration, 5 µl of sample was taken, and distilled water (995 µl) was added. The sample was then diluted by a factor of 10 and total protein

concentration was measured spectrophotometrically at 280 nm. The total protein concentration was used as a measure of the biomaterial in the experiments. For measurement of MDA concentration, homogenised sample (500 µl) was mixed with buffer A, 30% trichloroacetic acid, 0.75% 2-thiobarbituric acid, 0.5 M HCl and 0.02% butylated hydroxytoluene (500 ml), incubated at 90 °C for 30 min, then chilled on ice. n-Butanol (1.5 ml) was mixed with the sample, and the mixture was centrifuged at 6700 rcf for 10 min at 4 °C. The absorbance of the resulting chromophore was measured at 535 nm and 600 nm and the latter was subtracted from the former to correct for nonspecific turbidity. The concentration of MDA was calculated using an extinction coefficient of 156 mM⁻¹ cm⁻¹ (Ortega-Villasante et al., 2005). For statistical analysis, each concentration of MDA was divided by the total protein concentration of the corresponding sample.

Reactive oxygen species (ROS) assessment.

Assessment of ROS was performed by using the OxiSelect Intracellular ROS Assay KitTM (Cell Biolabs) measuring green fluorescence as described by (Petkovic et al., 2011b). DCFH-DA (2',7'-dichlorodihydrofluorescein diacetate), standards, H₂O₂ and TiO₂ suspension were prepared in cell media (PM). *T. thermophila* cells were first pretreated with 100 µM solution of DCHF-DA in the PM cell culture media for 60 min at 30 °C. Cells were then treated with 250 µM H₂O₂ and 0.1 and 100 µg/ml nano-TiO₂ particles or 0.1 and 100 µg/ml bulk-TiO₂ particles for 4 h. The DCF fluorescence intensity is proportional to the amount of ROS formed intracellularly. H₂O₂ is the principle ROS, responsible for the oxidation of DCFH-DA to DCF (LeBel et al., 1992). Negative (nontreated cells) and positive (H₂O₂ treated cells) controls were included in each experiment. For kinetic analysis of ROS formation the plates were maintained at 25 °C and the fluorescence intensity (480 nm excitation / 530 nm emission wavelengths) of the DCF formed was recorded every 5 min (for the first 30 min) and then every 30 min during the remainder of the 4 h incubation, using a Synergy H4 hybrid fluorescence plate reader (BioTrek). The statistical significance between treated groups and controls was determined by two-tailed Student's t-test

and $P < 0.05$ was considered as statistically significant. For each concentration of nano or bulk TiO_2 , three independent assays and two technical replicates were carried out.

Comet assay.

Different protocols and versions of Comet assay were used to assess the extent and the type of DNA damage as shown in Figure 1.

Comet assay with alkaline lysis *in vivo*.

The alkaline version of the Comet assay was performed by modifications of the original protocol (Lah et al., 2004). After exposure to TiO_2 particles in PM (as described in "Exposure Conditions, above), cells were harvested by 5 min centrifugation at 60 rcf washed with PM and resuspended in PM. To achieve a uniform background, rough microscope slides were coated with 400 μl of 0.5% normal melting point (NMP) agarose and were left to air dry overnight. Cells were mixed with 3.0% low melting point (LMP) agarose and spread over the slides as the second layer, giving a final concentration of 140 cells/ μl . After removing the cover glasses, the slides were covered with a third layer, of 300 μl of 3.5 % LMP agarose, to prevent escape of *T. thermophila* DNA during cell lysis and electrophoresis.

T. thermophila cells embedded in agarose were dipped in phosphate-saline buffer (PBS; 80 g NaCl, 8 g NaCl, 2 g KCl, 2 g KH_2PO_4 in 1 L doubly distilled H_2O at pH 7.2 to 7.4) for 20 min on ice and then washed twice with PBS. Slides were incubated overnight in lysis solution (30 mM NaOH, 1.2M NaCl, 1% (w/v) laurylsarcosine, 0.05% Triton X 100, 1% DMSO pH 12.4). The slides were rinsed three times for 20 min each in electrophoresis buffer (30 mM NaOH, 10 mM EDTA, pH 12.4) to remove lysis solution and to unwind the nuclear DNA. The samples were then subjected to electrophoresis for 20 min at 25 V and 300 mA in the same buffer. Following the electrophoresis the gels were neutralized in 400 mM Tris-HCl, pH 7.5 for 15 min. For visualization in a fluorescence microscope, the slides were stained with ethidium bromide (10 $\mu\text{g}/\text{ml}$) and 60 randomly selected nuclear images of each slide were acquired with an

epifluorescent microscope (Olympus BX50), using a BP 515-560 nm excitation filter and a barrier filter of LP 590 nm at 400 x magnification (Fig 2). Microscopic images of comets were captured by a digital camera (Hamamatsu Orca 2), connected to a computer. Detected comets were scored by Comet 5.0 Computer Software (Kinetic Imaging Ltd., 2001). The tail lengths and percentage of DNA in the comet's tails and heads were determined and further used to analyze the nuclear DNA damage.

Comet assay with alkaline lysis *in vitro*.

After culture growth in RM for 24 h in the dark at 25°C, cells were harvested by 5 min centrifugation at 60 rcf, washed with PM and resuspended in PM. The cells were embedded into 3.0 % low melting point agarose, the first and the third layer prepared as described above. Glass slides with embedded cells were then exposed to TiO₂ particles in PM for 1 h and then treated with TiO₂ nano and bulk particles (0.1 and 100 µg/ml) for 4 h. *T. thermophila* cells embedded in agarose were dipped in PBS for 20 min on ice and then washed twice with PBS. Glass slides were treated in alkaline lysis and all further steps were the same as described in the section “Comet assay with alkaline lysis *in vivo*” above.

Acellular Comet assay with alkaline lysis.

After culture growth in RM for 24 h in the dark at 25°C, cells were harvested by 5 min centrifugation at 60 rcf, then washed with PM and resuspended in PM. The cells were embedded into 3.0 % LMP agarose and the first and the third layers were prepared as described above. *T. thermophila* cells embedded on glass slides were dipped in PBS for 20 min on ice and then washed twice with PBS. Glass slides were treated after alkaline lysis and washed 3 times with PBS buffer for 10 min. One hour exposure of embedded nuclei to PM in the dark at 25°C was followed by the exposure to TiO₂ particles at two selected concentrations (TiO₂ nano- and bulk particles; 0.1 and 100 µg / ml) for 4 h, in the dark at 25 °C. After exposure, the glass slides were washed with electrophoresis buffer (EF buffer; 6 mL NaOH , 4 mL EDTA, 1990 mL MQ) and all further steps were the same as in section “Comet assay with alkaline lysis *in vivo*” above.

Comet assay with neutral lysis *in vivo* and *in vitro*.

Both *in vivo* and *in vitro* exposures to TiO₂ particles were tested as described in the sections “Comet assay with alkaline lysis *in vivo*” and “Comet assay with alkaline lysis *in vitro*” above. As a positive genotoxic control toxicant, 100µM methyl methanesulfonate (MMS) was used. For the Neutral Comet assay, a modification of the protocol by Wojewodzka et al. (2002) was used. The cell suspension was mixed with low melting point agarose (LMP agarose) at a final concentration of 0.75%. After the preparation of the third layer, the slides were left at 4 °C in the dark for 1-2 h in the lysing buffer which consisted of 2.5 M NaCl, 100 mM EDTA, 10 mM Tris-HCl, 1% *N*-lauroylsarcosine, pH 9.0. Immediately before use, 0.5% Triton X-100 and 10% dimethylsulfoxide (DMSO) were added to the buffer and mixed for 20 min. After 1 h of lysis, the slides were washed three times with the electrophoresis buffer (300mM sodium acetate, 100mM Tris-HCl, pH 8.3) and left in fresh buffer solution for 1 h, then placed in a horizontal gel electrophoresis unit filled with a fresh electrophoretic buffer. The slides were electrophoresed for 1 h at 14V (0.5 V/cm, 11–12 mA) at 8 °C.

Statistical analysis of Comet assay results.

The average percentage of tail DNA was compared in an incomplete four factor experimental design using analysis of variance (ANOVA). The first factor was “lysis” with two levels: alkaline and neutral; the second factor was “method” with four levels: *acellular* comet assay (only by alkaline lysis), *in vivo* Comet assay, *in vitro* comet assay and control. The third factor was the “size” of TiO₂ particles, either nano or bulk; and the fourth factor was concentration of TiO₂ particles, either: 0.1 or 100 µg/ml. The experiment was carried out in three biological replicates. At least 60 nuclei were examined in each replicate and the medians of percentage of tail DNA were calculated for each biological replication. The ANOVA calculations were made on the basis of the medians of percentage of tail DNA. The Duncan’s multiple

comparison test was used to determine the statistical significant differences between the treatments ($\alpha = 0.05$).

RESULTS

CHARACTERIZATION OF TiO₂ NANOPARTICLE SUSPENSIONS

The TEM revealed that TiO₂ nanoparticles were homogeneous in shape and size, with an aspect ratio of up to 1:5 between the diameter and length, forming elongated, spheroidal shapes. The transmission-electron diffraction pattern showed the TiO₂ to be in its anatase phase. BET analyses revealed the surface area to be between 190 and 290 m²/g and the average particle size to be 15 nm.

Dynamic light scattering analysis showed the average value of the hydrodynamic radius R_h , of TiO₂ nanoparticles suspended in test medium to be 820 nm. The average size of bulk TiO₂ could not be measured accurately with this approach because of the presence of larger agglomerates.

Zeta potentials of TiO₂ nanoparticle suspension (1000 µg/ml) were measured in the same medium used to expose cells, at pH 7.4. The value recorded was -15, which is equivalent to a suspension of incipient stability (Fig. 2).

CELLULAR FATTY ACID COMPOSITION

No significant differences have been found in membrane fatty acid profiles of *T. thermophila* after exposure to different concentrations of nano- or bulk TiO₂ at 25 °C after 4 h (Table 1). This suggests that TiO₂ particles have no effect on *T. thermophila* cell membranes.

LIPID PEROXIDATION

There were no differences in lipid peroxidation of analyzed *T. thermophila* samples after 0 and 4 h of incubation at 25 °C with nanoparticles, when compared to control cells. The average content of malondialdehyde in the control samples was 140 ± 23 nM of MDA per mg of protein.

REACTIVE OXYGEN SPECIES (ROS) PRODUCTION

In comparison to control cells, no ROS production was detected after 4 h of incubation with TiO₂ particles of any size at a concentration of 0.1 µg/ ml, however at 100 µg/ ml bulk-TiO₂, but not nano-TiO₂, significant elevation of intracellular ROS formation was detected (Fig. 3A).

To explore whether TiO₂ nanoparticles (0.1 and 100µg/ml) induced ROS formation not only at the end of exposure but also during the experiment, we measured the kinetics of their formation in *T. thermophila* cells in different time frames during 4 h of exposure (Fig. 3B). Comparison between treated groups and controls tested by two-tailed Student's t-test and $P < 0.05$, showed no statistically significant changes in ROS formation.

COMET ASSAY

Statistical analysis of the results obtained with a Comet assay after alkaline lysis indicated significant damage of DNA in *T. thermophila* in both *in vivo* and *in vitro* treatments with TiO₂ in comparison to control. This was independent of both the size and the concentration of particles (Fig. 4).

Statistical analysis of results of a Comet assay obtained with embedded nuclei (*acellular* exposure) also showed significant DNA damage at all TiO₂ exposure concentrations and sizes used, except for 100 µg/ml nano-TiO₂ concentration. Statistically significant differences, calculated using Duncan's multiple comparison test, between DNA damage in the two exposure concentrations of particles have been observed

in *acellular* Comet assays. A possible explanation for this is that nano- and bulk particles in suspensions aggregate more at higher concentrations (100 µg/ml) and this may hinder penetration into the gels.

Statistical analysis of Comet assays by neutral lysis showed that in cells treated with TiO₂ the average DNA tail length does not significantly differ from that in control cells, indicating no double strand breaks occur as a result of exposure to TiO₂ (Fig. 4). Double strand breaks did not occur in bulk- or in nano-TiO₂ treated cells and concentration and exposure type, namely in *in vivo* or *in vitro* experiments, failed to produce double strand breaks in DNA. When cells were treated with 100µM MMS, a reference positive control for double strand breaks, a statistically significant level of DNA damage was recorded.

Simultaneous performance of alkaline lysis and neutral lysis in this study indicates that single strand breaks are the main category of DNA damage caused by TiO₂ particles. No double strand breaks were observed (Fig. 4). The results of the *acellular* exposure to TiO₂ revealed the capacity of TiO₂ particles to produce extensive single strand breaks when interacting with embedded nuclei and implies that when TiO₂ particles are present in the medium during in a Comet assay they can interfere with DNA and give rise to false positive results and overestimates of actual genotoxicity.

JUST ACCEPTED

DISCUSSION

We report experimental evidence of TiO₂ particle interactions with DNA during the Comet assay that resulted in a positive test result. We studied the DNA damage sustained by *T. thermophila* incubated with TiO₂ bulk and nanoparticles and assessed by a Comet assay and we analysed cellular responses, including lipid peroxidation, ROS formation and membrane fatty acid profiles. The DNA was exposed to particles in three different exposure scenarios in order to assess whether nanoparticles could directly interact with DNA during the course of the assay —and thus produce a false positive result or an overestimate of the actual genotoxicity.

The ability of TiO₂-NPs to damage DNA has been shown in many studies (Gurr et al., 2005; Wang et al., 2007; Trouiller et al., 2009, see (Karlsson, 2010); see (Sathya et al., 2010)), but it has generally been rationalized as a consequence of oxidative stress. Our results showed that only TiO₂ bulk particles at 100 µg/ml cause significant ROS production, a result never observed with nano-particles. Other cellular markers such as membrane fatty acid profiles and lipid peroxidation, which could be also regarded as markers of cytotoxicity, remained unchanged compared with control cells. ROS elevation by bulk-TiO₂ particles (100 µg/ml) does not imply a higher degree of DNA damage and these results clearly indicate that oxidative stress is not a cause of the genotoxicity which was detected in our Comet assay study. Consequently, the recorded genotoxicity must be either independent of oxidative stress or a false positive result. Since literature data failed to report direct primary genotoxicity but rather genotoxicity driven by oxidative stress, the Comet assay results would appear to be false positives. There is only one published study in which, judging by an alkaline Comet assay, no genotoxicity was observed with nanoparticulate TiO₂ (Bhattacharya et al., 2009). We hypothesize that in this case, complete removal of the nanoparticles from the test system was achieved and the particles were not endocytosed to any significant degree. Potential causes of false positive results include particles which may remain in the test medium or particles which are present in food vacuoles or have been endocytosed. That intracellular particles can gain access

to DNA after lysis in the course of a Comet assay has been discussed by Stone et al., (2009) (Stone et al., 2009)(Stone et al., 2009) and by Karlsson (2010).

Our results agree with those in other reports. A review by Landsiedel *et al.* (Landsiedel et al., 2009) reported results of nanomaterials genotoxicity tests which were dependent upon the tests themselves. In an assessment of the genotoxicity of nanoparticles, in six studies the Ames test showed no genotoxicity, and this was associated with a barrier to penetration by the nanomaterials through the bacterial cell wall. In contrast, in 12 of 14 *in vitro* micronucleus assays 12 produced evidence of genotoxicity and in the Comet assay 14 of 19 studies showed nanomaterials to be highly genotoxic (Landsiedel et al., 2009). A partial explanation for these inconsistencies among the tests may be the fact that the Comet assay is the most sensitive of the assays but since different concentrations of nanoparticles were applied in the studies, this suggestion cannot be a complete explanation and other factors, such as direct interaction of NPs with DNA during the tests should be considered.

Based on the results of our study, presented here we suggest when the Comet assay is selected for assessment of genotoxicity of nanoparticles, pretesting of potential of nanoparticles to interact with DNA *post-festum* must be carried out. One means by which such interactions could be detected is use of the *acellular* Comet test. In addition, before settling on the Comet assay it is important to know whether to expect substantial amounts of intracellular nanoparticles which could interact with DNA while the test is proceeding.

Suspected genotoxicity should be confirmed by an independent assay or, at a minimum, with biomarkers indicating DNA repair, for example, mRNA expression of tumor suppressor gene p53 and its downstream regulated responsive genes (Petkovic et al., 2011b)), DNA deletions (Trouiller et al., 2009), inflammation (Trouiller et al., 2009, Grassian et al., 2007), or indications of oxidative stress status such as lipid peroxidation, or elevated levels of ROS (Gurr et al., 2005, Kang et al., 2008).

In isolation, the results of Comet assays are unreliable as a measure of nanoparticles' genotoxicity due to the possibility of false positives. In the future, the test protocol needs modifications in terms of exclusion or control of particle-assay interactions and combination with other oxidative stress markers. Only with such refinements will the Comet assay remain a test capable of reliably confirming or disproving genotoxicity.

CONCLUSIONS

- 1) Genotoxicity of TiO₂ nanoparticles was demonstrated when *T. thermophila* cells were incubated with nano-TiO₂ or bulk TiO₂ in a suspension (*in vivo* exposure), or embedded in gels (*in vitro* exposure) or when only embedded nuclei (*acellular* exposure) were exposed to nanoparticles (Fig. 1). Since positive Comet assay results were not accompanied by cytotoxicity markers such as lipid peroxidation, ROS formation or changes in composition of cell membranes, our Comet assay results appear to represent a false positive.
- 2) We suggest that in the future, pretesting of particle DNA interactions should be conducted in an *acellular* Comet assay and only the Comet assay results consistent with this pretesting should be accepted.
- 3) Data obtained from a Comet assay method alone are inadequate to support an assertion of an enhancement of the genotoxic potential of NPs. The genotoxic potential of NPs as obtained by a Comet assay should only be accepted when combined with evidence adduced by properly selected oxidative stress biomarkers.

Declaration of interest

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

Acknowledgements

This study was supported by Slovenian Research Agency: Projects no. 1000-07-310129 and no. J1—4109.

We thank Professor Bill Milne for critical reading of the manuscript.

REFERENCES

- ASHARANI, P. V., LOW KAH MUN, G., HANDE, M. P. & VALIYAVEETIL, S. 2009. Cytotoxicity and genotoxicity of silver nanoparticles in human cells. *ACS Nano*, 3, 279-90.
- BANASIK, A., LANKOFF, A., PISKULAK, A., ADAMOWSKA, K., LISOWSKA, H. & WOJCIK, A. 2005. Aluminum-induced micronuclei and apoptosis in human peripheral-blood lymphocytes treated during different phases of the cell cycle. *Environ Toxicol*, 20, 402-6.
- BHATTACHARYA, K., DAVOREN, M., BOERTZ, J., SCHINS, R. P., HOFFMANN, E. & DOPP, E. 2009. Titanium dioxide nanoparticles induce oxidative stress and DNA-adduct formation but not DNA-breakage in human lung cells. *Part Fibre Toxicol*, 6, 17.

- CHEN, M. & VON MIKECZ, A. 2005. Formation of nucleoplasmic protein aggregates impairs nuclear function in response to SiO₂ nanoparticles. *Exp Cell Res*, 305, 51-62.
- DHAWAN, A., BAJPAYEE, M. & PARMAR, D. 2009. Comet assay: a reliable tool for the assessment of DNA damage in different models. *Cell Biol Toxicol*, 25, 5-32.
- DIONISI, F., GOLAY, P. A., ELLI, M. & FAY, L. B. 1999. Stability of cyclopropane and conjugated linoleic acids during fatty acid quantification in lactic acid bacteria. *Lipids*, 34, 1107-15.
- DONALDSON, K., POLAND, C. A. & SCHINS, R. P. 2010. Possible genotoxic mechanisms of nanoparticles: criteria for improved test strategies. *Nanotoxicology*, 4, 414-20.
- FRANKEL, J. 2000. Cell biology of *Tetrahymena thermophila*. *Methods Cell Biol*, 62, 27-125.
- GONZALEZ, L., SANDERSON, B. J. & KIRSCH-VOLDERS, M. 2011. Adaptations of the in vitro MN assay for the genotoxicity assessment of nanomaterials. *Mutagenesis*, 26, 185-91.
- GRASSIAN, V. H., O'SHAUGHNESSY P, T., ADAMCAKOVA-DODD, A., PETTIBONE, J. M. & THORNE, P. S. 2007. Inhalation exposure study of titanium dioxide nanoparticles with a primary particle size of 2 to 5 nm. *Environ Health Perspect*, 115, 397-402.
- GREIM, H. & NORPPA, H. 2010. Genotoxicity testing of nanomaterials--conclusions. *Nanotoxicology*, 4, 421-4.
- GURR, J. R., WANG, A. S., CHEN, C. H. & JAN, K. Y. 2005. Ultrafine titanium dioxide particles in the absence of photoactivation can induce oxidative damage to human bronchial epithelial cells. *Toxicology*, 213, 66-73.
- GUTIÉRREZ, J. C., MARTÍN-GONZÁLEZ, A., DIAZ, S. & ORTEGA, R. 2003. Ciliates as a potential source of cellular and molecular biomarkers/biosensors for heavy metal pollution. *European Journal of Protistology*, 39, 461-467.
- HOLBROOK, R. D., MURPHY, K. E., MORROW, J. B. & COLE, K. D. 2008. Trophic transfer of nanoparticles in a simplified invertebrate food web. *Nat Nanotechnol*, 3, 352-5.

- KAHRU, A., DUBOURGUIER, H. C., BLINOVA, I., IVASK, A. & KASEMETS, K. 2008. Biotests and Biosensors for Ecotoxicology of Metal Oxide Nanoparticles: A Minireview. *Sensors*, 8, 5153-5170.
- KANG, S. J., KIM, B. M., LEE, Y. J. & CHUNG, H. W. 2008. Titanium dioxide nanoparticles trigger p53-mediated damage response in peripheral blood lymphocytes. *Environ Mol Mutagen*, 49, 399-405.
- KARLSSON, H. L. 2010. The comet assay in nanotoxicology research. *Anal Bioanal Chem*, 398, 651-66.
- LAH, B., MALOVRH, S., NARAT, M., CEPELJNIK, T. & MARINSEK-LOGAR, R. 2004. Detection and quantification of genotoxicity in wastewater-treated *Tetrahymena thermophila* using the comet assay. *Environ Toxicol*, 19, 545-53.
- LANDSIEDEL, R., KAPP, M. D., SCHULZ, M., WIENCH, K. & OESCH, F. 2009. Genotoxicity investigations on nanomaterials: methods, preparation and characterization of test material, potential artifacts and limitations--many questions, some answers. *Mutat Res*, 681, 241-58.
- LANDSIEDEL, R., MA-HOCK, L., VAN RAVENZWAAY, B., SCHULZ, M., WIENCH, K., CHAMP, S., SCHULTE, S., WOHLLEBEN, W. & OESCH, F. 2010. Gene toxicity studies on titanium dioxide and zinc oxide nanomaterials used for UV-protection in cosmetic formulations. *Nanotoxicology*, 4, 364-81.
- LEBEL, C. P., ISCHIROPOULOS, H. & BONDY, S. C. 1992. Evaluation of the probe 2',7'-dichlorofluorescein as an indicator of reactive oxygen species formation and oxidative stress. *Chem Res Toxicol*, 5, 227-31.
- LIN, M. H., HSU, T. S., YANG, P. M., TSAI, M. Y., PERNG, T. P. & LIN, L. Y. 2009. Comparison of organic and inorganic germanium compounds in cellular radiosensitivity and preparation of germanium nanoparticles as a radiosensitizer. *Int J Radiat Biol*, 85, 214-26.
- NEL, A., XIA, T., MADLER, L. & LI, N. 2006. Toxic potential of materials at the nanolevel. *Science*, 311, 622-7.

- ORTEGA-VILLASANTE, C., RELLAN-ALVAREZ, R., DEL CAMPO, F. F., CARPENA-RUIZ, R. O. & HERNANDEZ, L. E. 2005. Cellular damage induced by cadmium and mercury in *Medicago sativa*. *J Exp Bot*, 56, 2239-51.
- PETKOVIC, J., KUZMA, T., RADE, K., NOVAK, S. & FILIPIC, M. 2011a. Pre-irradiation of anatase TiO₂ particles with UV enhances their cytotoxic and genotoxic potential in human hepatoma HepG2 cells. *J Hazard Mater*, 196, 145-52.
- PETKOVIC, J., ZEGURA, B., STEVANOVIC, M., DRNOVSEK, N., USKOKOVIC, D., NOVAK, S. & FILIPIC, M. 2011b. DNA damage and alterations in expression of DNA damage responsive genes induced by TiO₂ nanoparticles in human hepatoma HepG2 cells. *Nanotoxicology*, 5, 341-53.
- SATHYA, T. N., VARDHINI, N. V. & BALAKRISHNAMURTHY, P. 2010. REVOLUTION OF 'NANO' IN IN-VITRO GENETIC TOXICOLOGY. *Journal of Cell and Tissue Research* Vol. 10, 2389-2396
- SAUVANT, M. P., PEPIN, D. & PICCINNI, E. 1999. *Tetrahymena pyriformis*: a tool for toxicological studies. A review. *Chemosphere*, 38, 1631-69.
- SCHULTZ, T. W. 1997. Influence of the energy relationship of organic compounds on toxicity to the cladoceran *Daphnia magna* and the fish *Pimephales promelas*. *Ecotoxicol Environ Saf*, 38, 336-8.
- STONE, V., JOHNSTON, H. & SCHINS, R. P. 2009. Development of in vitro systems for nanotoxicology: methodological considerations. *Crit Rev Toxicol*, 39, 613-26.
- TICE, R. R., AGURELL, E., ANDERSON, D., BURLINSON, B., HARTMANN, A., KOBAYASHI, H., MIYAMAE, Y., ROJAS, E., RYU, J. C. & SASAKI, Y. F. 2000. Single cell gel/comet assay: guidelines for in vitro and in vivo genetic toxicology testing. *Environ Mol Mutagen*, 35, 206-21.
- TROUILLER, B., RELIENE, R., WESTBROOK, A., SOLAIMANI, P. & SCHIESTL, R. H. 2009. Titanium dioxide nanoparticles induce DNA damage and genetic instability in vivo in mice. *Cancer Res*, 69, 8784-9.

WARHEIT, D. B. & DONNER, E. M. 2010. Rationale of genotoxicity testing of nanomaterials: regulatory requirements and appropriateness of available OECD test guidelines. *Nanotoxicology*, 4, 409-13.

JUST ACCEPTED

Table and figures:

Table 1. Average percentage composition of membrane fatty acid samples from *T. thermophila* exposed to TiO₂ particles at 25°C after 4 h. The data are presented as total sums of various fatty acids of lipid extracted from the three independent cultures. Percentages are expressed as means ± standard error (SE). B = bulk concentration, NP = nanoparticle concentration, FA = fatty acid.

JUST ACCEPTED

Figure 1. The protocols and types of exposure used in our genotoxicity study. These indicate where and when the bulk- and nano-TiO₂ particles may remain in close proximity to nuclei in the final steps of the Comet assay, leading to an overestimate of genotoxicity and type of DNA damage (DSS=double DNA strand breaks, SSB=single DNA strand breaks).

Figure 2. Zeta potentials of TiO₂ nanoparticle suspensions (1000 µg/ml) measured in the PM medium and used in experimental exposures.

Figure 3. Induction of ROS formation in *T. thermophila* cells. (A) treated with H₂O₂ (250 µM), nano-TiO₂ particles (NP, 0.0, 0.1 and 100µg/ml), bulk-TiO₂ particles (0.1 and 100µg/ml) and presented as a relative increase of DCF fluorescence after 4 h of exposure to TiO₂ particles. Each bar is represented as a mean ± standard error (SE) of three independent experiments. (B) Kinetics of ROS formation during exposure for 4 h to TiO₂ NPs (0.1 and 100 µg/ml). Each point represents the mean of 6 replicates ± standard error (SE).

Figure 4. Results of the Comet assay experiment. *T. thermophila* was treated with nanoparticles (NP) or bulk TiO₂ particles (B) at two different concentrations (1, 0.1 µg/ml and 2, 100 µg/ml). Three different exposure conditions (*acellular, in vitro, in vivo*) were applied and two different protocols of Comet assay (alkaline lysis and neutral lysis) were used.

Table 1.

Particle type Particle concentration (µg/ml)	Bulk TiO ₂ (B)						Nano TiO ₂ (NP)					
	0	0,1	1	10	100	1000	0	0,1	1	10	100	1000
Straight chain saturated FA (%)	37.2±0.9	36.2±0.4	37.1±0.8	38.3±0.1	36.0±0.6	37.4±0.4	31.0±0.4	31.0±0.5	30.8±0.8	31.3±0.1	31.5±0.5	32.4±0.2
Unsaturated FA (%)	49.6±1.0	50.1±0.4	50.0±1.1	48.3±0.3	50.6±0.8	49.4±0.4	52.0±0.4	51.6±0.6	52.5±0.6	51.7±0.6	51.5±0.7	50.1±0.4

monounsaturated FA (1x) (%)	22.4±0.6	21.3±1.3	21.2±0.8	22.6±1.5	21.4±0.5	23.7±1.2	22.3±0.5	22.3±0.5	21.7±1.0	22.7±0.3	21.8±0.6	22.1±0.3
polyunsaturated FA (2x) (%)	8.3±.2	8.6±.2	8.5±.2	8.0±.1	8.6±.2	8.1±.1	8.8±.0	8.3±.5	8.7±.1	8.7±.2	8.7±.1	8.4±.1
polyunsaturated FA (3x) (%)	18.9±1.4	20.3±1.3	20.2±1.7	17.7±1.2	20.6±1.1	17.7±1.3	21.0±0.9	21.1±0.3	22.0±1.0	20.3±0.7	21.1±0.4	19.6±0.6
Saturated iso & anteiso branched FA (%)	5.6±.3	5.4±.1	5.6±.2	5.9±.1	5.4±.1	5.8±.2	7.3±.4	7.5±.1	7.6±.1	7.7±.2	7.5±.2	7.8±.3
saturated iso FA (%)	5.0±.3	4.8±.1	5.0±.2	5.2±.1	4.8±.1	5.1±.1	6.5±.1	6.4±.1	6.5±.0	6.6±.2	6.5±.1	6.7±.3
saturated anteiso FA (%)	0.6±.0	0.6±.0	0.6±.0	0.6±.0	0.7±.1	0.7±.1	0.7±.4	1.1±.0	1.1±.0	1.1±.0	1.1±.0	1.1±.0
Average number of C-atoms in membrane FA	14.7	14.9	15.0	14.9	14.9	15.0	14.7	14.7	14.9	15.0	14.7	14.6

JUST ACCEPTED




Figure 1.

JUST ACCEPTED






Figure 2.



A



Figures 3 A and B.




Figure 4.

Figure 4. Results of the Comet assay experiment. *T. thermophila* was treated with nanoparticles (NP) or bulk TiO₂ particles (B) at two different concentrations (1, 0.1 μg/ml and 2, 100 μg/ml). Three different exposure conditions (*acellular*, *in vitro*, *in vivo*) were applied and two different protocols of Comet assay (alkaline lysis and neutral lysis) were used. There is no statistically significant difference between averages indicated with the same letter (a,b,c,...., f).






Figure 2.



A



B
Figures 3 A and B.




Figure 4.